481

by Susy Saadah

Submission date: 24-Nov-2023 11:54AM (UTC+0700)
Submission ID: 2095099211
File name: 481-Article_Text-1675-1-2-20231110.doc (87.5K)
Word count: 2231
Character count: 12942

SANITAS: JURNAL TEKNOLOGI DAN SENI KESEHATAN
ISSN : 1978-8843 (*PRINT*) / 2615-8647 (*ONLINE*)
DOI: 10.36525/sanitas.2023.i1
Vol. 14 No. 1: xx - xx

COMPARISON OF ANTIOXIDANT ACTIVITY OF CRUDE EXTRACT AND GEL PREPARATION FROM Alstonia scholaris L. LEAF EXTRACT

ABSTRACT

Alstonia scholaris L. contains several compounds, which has been proven to be potential as skin protection by formulating in gel preparation. The aims of this study were to obtained the antioxidant activity of crude extract from leaf part of Alstonia scholaris L. and gel preparation contain Alstonia scholaris L. leaf extract. Extract was prepared using destilled methanol by maceration method. The gel preparation was developed with extract (1% w/v) in carbomer base (1,67% w/v). The chemical compound was evaluated qualitatively used some reagent. The antioxidant assays performed by DPPH (2,2-diphenyl-1-picrylhydrazyl) method. The result shown that methanolic extract of Alstonia scholaris leaf was found had flavonoids, tannins, phenols, alkaloids, saponins, and steroids compound. The methanolic crude extract and gel preparation had strong antioxidant activity with IC50 value was found to be 69,502 ppm and 86,832 ppm, respectively. From the result we conclude that the methanolic crude extract of Alstonia scholaris leaf sole has strong scavenging free radical activity even formulated in gel preparation but has decreased IC50 value.

Keywords: Alstonia scholaris; antioxidant; DPPH; gel; carbomer

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non-Commercial-Share Alike 4.0 license, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.
©2023 Sanitas

DOI: 10.36525/sanitas.2023.i1
Vol. 14 No. 1: xx - xx

INTRODUCTION

Alstonia scholaris L. is one of the plants that is widely used by the community as a medicinal plant (Baliga, M.S, 2012). The most frequently isolated active compounds from this genus are Monoterpenoid Indole Alkaloids (MIAs) (M. xue Zhao et al, 2023). Apart from that, the phenolic compounds, flavonoids and proanthocyanidins were proven to be found in the leaf, follicles and latex part of this plant with the highest content in the leaf with phenol levels of 49.66 ± 1.52 mg GAE/g, flavonoid levels of 97.33 ±1.52 mg QE/g and proanthocyanidin levels 99.33±1.52 mg CE/g (Ganjewala and Gupta, 2013).

Research was proven that *Alstonia scholaris* L. has antioxidant, antibacterial, antihyperuricemia and anticancer activity (Manjeshwar, 2010; Ganjewala and Gupta, 2013; Hu BY, et al, 2022). Other research also showed that *Alstonia scholaris* L. has antitussive, expectorant, anti-asthmatic, anti-inflammatory and analgesic activity (Shang JH et al, 2010; Shang JH et al 2010).

However, several studies show that there are differences of chemical active compounds in *Alstonia scholaris* L. in several places where they grow, causing differences in the pharmacological activity of these plants, including activity as antioxidants (Saxena N et al, 2012; Kaushik P et al, 2011; Arulmozhi S et al, 2010; Ramachandra YL, et al, 2012). On the other hand, the formulation can cause some differences in pharmacological activity from before it was formulated due to the presence of excipients (Sousa R, 2021). Therefore, in this research was carried out the differences of antioxidant activity of *Alstonia scholaris* L. ethanolic extract and after formulation in gel preparations using native Indonesian plants.

MATERIAL & METHODS

Collecting samples and simplicia process

Alstonia scholaris L. leaf were collected from the ex-MTQ area of Jambi City, Jambi Province and determined at the Plant Taxonomy Laboratory, Biology Department, Faculty of Mathematics and Natural Sciences, Padjadjaran University. Fresh leaf were taken, wet sorted then dried used oven at 45°C. The simplicia was powdered with a grinder and the

DOI: 10.36525/sanitas.2023.i1
Vol. 14 No. 1: xx - xx

percentage of yield was calculated by comparing the weight of simplicia to the initial weight of the sample.

Preparation of Extract

The extract was made using methanol distillate as solvent with maceration method for 3x24 hours with occasional stirring. the ratio of simplicia and solvent were 1:10. The maserate was collected and the solvent was evaporated using a rotary evaporator at a temperature of 50°C until a thick extract was obtained. the percent of yield was calculated by comparing the weight of the extract obtained to the weight of the initial simplicia powder used.

Non-Specific and Specific Parameters, and Screening of Phytochemical

The methanolic extract of *Alstonia scholaris* L was determined non-specific parameters (water content and ash content) and specific parameters (identity and organoleptics). The extracts were identified chemical compounds of flavonoids, alkaloids, saponins, steroids, tannins and phenolics.

Gel Preparation

Formula of gel preparation was an optimal gel formula with extract concentration of 1%; carbopol 1.67%; glycerin 14.83%; methyl paraben 0.18%; propyl paraben 0.02%; NaOH 0.7%; and distilled water ad 100%. Gel preparation was expanded with carbopol in distilled water for 24 hours. Next, in a separate container, mix methyl paraben, propyl paraben, and glycerin while heating. The extract was added to the mixture and homogenized using homogenizer for 20 minutes. The expanded carbopol was mixed into the solution with stirred. Add NaOH and add distilled water, stir until homogeneous.

Antioxidant Activity

The antioxidant test was carried out using the DPPH (1,1-diphyenyl-2-picrylhydrazyl) method based on research by Zaky, et al (2021) with slight modifications. A series of antioxidant tests are carried out as follows:

a. Preparation of blank solution

   10 mL of 0,05 mM DPPH solution in methanol pro analysis was made and incubated in dark room. The absorbance of blank solution was measured by spectrophotometry method at 517 nm.

DOI: 10.36525/sanitas.2023.i1
Vol. 14 No. 1: xx - xx

b. Preparation of ascorbic acid solution

Ascorbic acid solution was made at a concentration of 1000 ppm in methanol pro analysis and diluted to 10; 20; 30; 40 and 50 ppm as positive control. Then, 3.8 mL of DPPH blank solution was homogenized with 0.2 mL various concentration of ascorbic acid solution, incubated in the dark room for 30 minutes then measured at a wavelength of 517 nm using a UV-Vis spectrophotometer.

c. Preparation of sample solution

Sample solution was made contained 10,000 ppm extract and diluted to a concentration of 400; 600; 800; 1000; and 2000 ppm. 3.8 mL of sample solution was added to 0.2 mL of DPPH blank solution. The solution was homogenized, stored in a dark place and the absorbance was measured at a wavelength of 517 nm using a UV-Vis spectrophotometer.

d. Determination of percent of inhibition and IC_{50} value

Percent of inhibition was calculated using this following formula:

Percent of inhibition=(blank absorbance-sample absorbance)/blank absorbancex100%. Then, linear regression equation was created between concentration (x) and percent inhibition (y). The IC_{50} value can be find as the x value from the regression equation with y=50.

RESULT & DISCUSSION

Plant determination was carried out with the result that was Alstonia scholaris (L.) R. Br from the Apocynaceae family. The results of simplicia was 4.388 kg of from 5,3 kg of sample. The results of the percent of simplicial and extract yield are presented in table 1.

Tablel 1. Percent of yield

Sample	Percent of yield (%)
Simplicia	82,79
Crude extract	5,7

Determination of plants is the initial stage of research which aims to determine and ensure the viability of the plants in this research. The samples used were fresh pulai leaf which is

DOI: 10.36525/sanitas.2023.i1
Vol. 14 No. 1: xx - xx

hoped increase the economic value of these plant. Pulai leaf simplicia was extracted by maceration method, therefore components that cannot tolerate heating are still present in the extract (Taslim, 2020).

Crude extract was determined the specific, non-specific parameters and phytochemical screening, shown in table 2 and table 3.

Table 2. Nonspecific and specific parameter of crude extract

Parameters	Result
Nonspecific parameters	
Water content	1,8%
ash content	1,87%
Specific parameters	
Identity	
Extract	*Alstonia scholaris* L. methanolic Leaf extract
Botanical Name	*Alstonia scholaris* L.
Part of plant	Folium (leaf)
Indonesian Name	Pulai, Pule
Organoleptics	
Smell	Typical Pulai Extract Smell
Type of extract	Dry extract
Flavor	Bitter
Color	Blackish Green

Based on the screening results, pulai leaf contain secondary metabolites, such as flavonoids, phenolics, tannins, alkaloids, saponins and steroids. The orange color from the identification of flavonoids was due to magnesium powder can reduction reaction with flavonoid compounds resulting in a change in the color of the extract. Because of reaction with FeCl3, crude extract contain phenolic and tannins compound become blue or blackish green.

The identification reaction of alkaloids produced precipitate because the nitrogen atom in alkaloid structure with a lone pair of electrons substitutes the iodine ion in Dragendorf reagent covalently.

Furthermore, because of saponin has hydrophilic and hydrophobic groups, the extract containing saponin compounds generates foam and stable when 2 N HCl was added. HCl

DOI: 10.36525/sanitas.2023.i1
Vol. 14 No. 1: xx - xx

can increase the polarity of saponin caused by the polar (hydrophilic) groups to face outward and the non-polar (hydrophobic) groups to face inward, resulting in the formation of a micelle. *Alstonia scholaris* L. leaf extract also contains steroid compounds which are characterized by the formation of a green or blue color due to oxidation through the formation of conjugated double bonds. These results were accordance with several studies, one was conducted by Kalaria, et al, 2012 shown that *Alstonia scholaris* L. leaf extract has flavonoids, phenolics, tannins, alkaloids, saponins and steroids (Putri, et al, 2020; Kalaria P, et al, 2012).

Table 3. Phytochemical screening of crude extract

Phytochemical compounds	Reagents	Result
Flavonoids	Mg powder+HCl	positive (orange color was formed)
Tannins	FeCl3	Positive (a blackish green/blackish blue color was formed)
Alkaloids	Dragendorf	positive (precipitated)
Saponins	Aquadest, HCl	positive (foams was formed)
Steroids	Anhydrate acetic acid+H2SO4	positive (blue color was formed)

Flavonoid molecules are the most compound that function as natural antioxidants. As antioxidants, flavonoids can provide radical molecules with hydrogen atoms. Nonetheless, the quantity and orientation of the OH groups determine how potently flavonoids exhibit antioxidant activity. The antioxidant activity of flavonoids increases with their hydroxyl group count.

The DPPH method was selected for the antioxidant activity test because imple, fast, and does not require a lot of reagents. With one unpaired of electron, DPPH is a radical molecule which gives it a purple color. According to Zaky (2021), discoloration from deep purple to clearer liquid indicates the formation of an electron pair (Zaky, M, 2021).

The results of determining the antioxidant activity of crude extracts and gel preparation shown that both have strong antioxidant activity. However, it was seen that the extract in

DOI: 10.36525/sanitas.2023.i1
Vol. 14 No. 1: xx - xx

gel preparation decrease in antioxidant activity as indicated by an increase in the IC50 value. The IC50 value is the sample concentration that can ward off 50% of free radicals. A smaller IC50 value indicates better antioxidant activity. The IC50 value of the sample (*Alstonia scholaris* L crude extract and *Alstonia scholaris* L extract in gel preparation) was very strong antioxidant activity, was still far from the IC50 value of the positive control (ascorbic acid).

CONCLUSION

The antioxidant activity of methanol extract of *Alstonia scholaris* L leaf extract and gel preparation containing this extract was strong category of antioxidant activity. However, there was a decrease in the value of its antioxidant activity after it was formulated in gel preparation.

REFERENCE

Arulmozhi S, Mazumder PM, Narayanan LS, Thakurdesai PA. In vitro antioxidant and free radical scavenging activity of fractions from *Alstonia scholaris* Linn.R.Br. *Int J Pharm Tech Res*. 2010;2(1):18–25.

Baliga, M.S. Review of the phytochemical, pharmacological and toxicological properties of *Alstonia Scholaris* Linn. R. Br (Saptaparna). *Chin. J. Integr. Med.* (2012). https://doi.org/10.1007/s11655-011-0947-0.

Ganjewala D, Gupta AK. Study on Phytochemical Composition, Antibacterial and Antioxidant Properties of Different Parts of *Alstonia scholaris* Linn. Adv Pharm Bull. 2013;3(2):379-84. doi: 10.5681/apb.2013.061. Epub 2013 Aug 20. PMID: 24312864; PMCID: PMC3848240.

Hu BY, Zhao YL, Ma DY, Xiang ML, Zhao LX, Luo XD. Anti-hyperuricemic bioactivity of *Alstonia scholaris* and its bioactive triterpenoids in vivo and in vitro. J Ethnopharmacol. 2022 May 23;290:115049. doi: 10.1016/j.jep.2022.115049. Epub 2022 Feb 9. PMID: 35150817.

Kalaria P, Gheewala P, Chakraborty M, Kamath J. A phytopharmacological review of *Alstonia scholaris*: A panoramic herbal medicine. Int J Res Ayurveda Pharm. 2012;3(3):367-371.

Kaushik P, Kaushik D, Sharma N, Rana AC. *Alstonia scholaris*: It's phytochemistry and pharmacology. *Chron Young Sci*. 2011;2(2):71–8.

Manjeshwar Shrinath Baliga. *Alstonia scholaris* Linn R Br in the treatment and prevention of cancer: past, present, and future. Integr Cancer Ther. 2010 Sep;9(3):261-9. doi: 10.1177/1534735410376068. Epub 2010 Aug 11. PMID: 20702494.

M. xue Zhao *et al*., "Traditional uses, chemical composition and pharmacological activities of *Alstonia* R. Br. (Apocynaceae): A review," *Arabian Journal of Chemistry*, vol. 16, no. 8. Elsevier B.V., Aug. 01, 2023. doi: 10.1016/j.arabjc.2023.104857.

Putri DM, Lubis SS. Skrining Fitokimia Ekstrak Etil Asetat Daun Kalayu (Erioglossum rubiginosum (Roxb.) Blum). Amina. 2020;2(3):120-125.

Ramachandra YL, Ashajyothi C, Padmalatha RS. Antioxidant activity of *Alstonia scholaris* extracts containing flavonoids and phenolic compounds. *Int J Pharm Pharma Sci*. 2012;4(3):424–6.

Saxena N, Shrivastava PN, Saxena RC. Preliminary physcio-phytochemical study of stem bark of *Alstonia scholaris* (L) RBR-A medicinal plant. *Int J Pharma Sci Res*. 2012;3(4):1071–5.

Shang JH, Cai XH, Feng T, Zhao YL, Wang JK, Zhang LY, Yan M, Luo XD. Pharmacological evaluation of *Alstonia scholaris*: anti-inflammatory and analgesic effects. J Ethnopharmacol. 2010 May 27;129(2):174-81. doi: 10.1016/j.jep.2010.02.011. Epub 2010 Feb 26. PMID: 20219658.

Shang JH, Cai XH, Zhao YL, Feng T, Luo XD. Pharmacological evaluation of *Alstonia scholaris*: anti-tussive, anti-asthmatic and expectorant activities. J Ethnopharmacol. 2010 Jun 16;129(3):293-8. doi: 10.1016/j.jep.2010.03.029. Epub 2010 Apr 8. PMID: 20381600.

Sousa R, Figueirinha A, Batista MT, Pina ME. Formulation Effects in the Antioxidant Activity of Extract from the Leaves of *Cymbopogon citratus*(DC) Stapf. Molecules. 2021 Jul 27;26(15):4518. doi: 10.3390/molecules26154518. PMID: 34361669; PMCID: PMC8348009.

Taslim T PR. Analisis Daya Antioksidan Senyawa Metabolit Sekunder Ekstrak Etanol Daun Surian (*Toona sinensis*(Juss) M.Roem).J Akad Farm Pray. 2020;5((2):19-28).

Zaky M, Rusdiana N, Darmawati A. Formulasi Dan Evaluasi Fisik Sediaan Gel Antioksidan Ekstrak Etanol 70% Daun Belimbing Wuluh (*Averrhoa Bilimbi* L.) Menggunakan Metode Dpph. J Farmagazine. 2021;8(2):26.

481

ORIGINALITY REPORT

20%	19%	10%	8%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	research.amanote.com Internet Source	3%
2	sanitas.e-journal.id Internet Source	3%
3	Submitted to Badan PPSDM Kesehatan Kementerian Kesehatan Student Paper	3%
4	jurnal.unimed.ac.id Internet Source	1%
5	jurnal.batan.go.id Internet Source	1%
6	www.myfoodresearch.com Internet Source	1%
7	Submitted to Universitas Brawijaya Student Paper	1%
8	oamjms.eu Internet Source	1%
9	journal.ubpkarawang.ac.id Internet Source	1%

10	innovareacademics.in Internet Source	1 %
11	Marlina, Salman, H Filza, E Nur, A N Poppy. "Antioxidant serum gel formulation with a combination of secretome from mesenchymal stem cells and rosemary oil", IOP Conference Series: Earth and Environmental Science, 2023 Publication	1 %
12	"Medicinal Plants - Recent Advances in Research and Development", Springer Science and Business Media LLC, 2016 Publication	<1 %
13	www.sciencegate.app Internet Source	<1 %
14	dspace.cus.ac.in Internet Source	<1 %
15	text-id.123dok.com Internet Source	<1 %
16	www.ncbi.nlm.nih.gov Internet Source	<1 %
17	Mi-xue Zhao, Jing Cai, Ying Yang, Jian Xu, Wen-yuan Liu, Toshihiro Akihisa, Wei Li, Takashi Kikuchi, Feng Feng, Jie Zhang. "Traditional uses, chemical composition and pharmacological activities of Alstonia R. Br.	<1 %

(Apocynaceae): A review", Arabian Journal of Chemistry, 2023
Publication

18	Rui Guo, Jian-Hua Shang, Rui-Han Ye, Yun-Li Zhao, Xiao-Dong Luo. "Pharmacological investigation of indole alkaloids from Alstonia scholaris against chronic glomerulonephritis", Phytomedicine, 2023 Publication	<1 %
19	www.researchgate.net Internet Source	<1 %
20	Elizabeth Elliott, François Chassagne, Agnès Aubouy, Eric Deharo et al. "Forest Fevers: traditional treatment of malaria in the southern lowlands of Laos", Journal of Ethnopharmacology, 2020 Publication	<1 %

Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	On		